


The groups included that have subsidiary status are

-Cl	Chloro	-NO ₂	Nitro	C=C	Double Bond
-Br	Bromo	-C≡N	Cyano	C≡C	Triple Bond
-I	Iodo	-OR	Alkoxy		Aromatic

The experiment presents all the chief chemical and spectroscopic methods of determining the main functional groups, and it includes methods for verifying the presence of the subsidiary functional groups as well. It will usually not be necessary to determine the presence of the subsidiary functional groups to identify the unknown compound correctly. Every piece of information helps the identification, however, and if these groups can be detected easily, you should not hesitate to determine them. Finally, complex bifunctional compounds are generally avoided in this experiment; only a few are included.

HOW TO PROCEED

Fortunately, we can detail a fairly straightforward procedure for determining all the necessary pieces of information. This procedure consists of the following steps:

Part One: Chemical Classification

1. Preliminary classification by physical state, color, and odor
2. Melting-point or boiling-point determination; other physical data
3. Purification, if necessary
4. Determination of solubility behavior in water and in acids and bases
5. Simple preliminary tests: Beilstein, ignition (combustion)
6. Application of relevant chemical classification tests

Part Two: Spectroscopy

7. Determination of infrared and NMR spectra

Part Three: Optional Procedures

8. Elemental analysis, if necessary
9. Preparation of derivatives
10. Confirmation of identity

Each of these steps is discussed briefly in the following sections.

PRELIMINARY CLASSIFICATION

Note the physical characteristics of the unknown, including its color, its odor, and its physical state (liquid, solid, crystalline form). Many compounds have characteristic colors or odors, or they crystallize with a specific crystal structure. This information can often be found in a handbook and can be checked later. Compounds with a high degree of conjugation are frequently yellow to red. Amines often have a fishlike odor. Esters have a pleasant fruity or floral odor. Acids have a sharp and pungent odor. A part of the training of every good chemist includes cultivating the ability to recognize familiar or typical odors. As a note of caution, many compounds have distinctly unpleasant or nauseating odors. Some

Experiment 51 Identification of Unknowns


Qualitative organic analysis, the identification and characterization of unknown compounds, is an important part of organic chemistry. Every chemist must learn the appropriate methods for establishing the identity of a compound. In this experiment, you will be issued an unknown compound, and will be asked to identify it through chemical and spectroscopic methods. Your instructor may give you a general unknown or a specific unknown. With a **general unknown**, you must first determine the class of compound to which the unknown belongs, that is, identify its main functional group; then you must determine the specific compound in that class that corresponds to the unknown. With a **specific unknown**, you will know the class of compound (ketone, alcohol, amine, and so on) in advance, and it will be necessary to determine only whatever specific member of that class was issued to you as an unknown. This experiment is designed so that the instructor can issue several general unknowns or as many as six successive specific unknowns, each having a different main functional group.

Although there are well over a million organic compounds that an organic chemist might be called on to identify, the scope of this experiment is necessarily limited. In this textbook, just over 300 compounds are included in the tables of possible unknowns given for the experiment (see Appendix 1). Your instructor may wish to expand the list of possible unknowns, however. In such a case, you will have to consult more extensive tables, such as those found in the work compiled by Rappoport (see References). In addition, the experiment is restricted to include only seven important functional groups:

Aldehydes	Amines
Ketones	Alcohols
Carboxylic acids	Esters
Phenols	

Even though this list of functional groups omits some of the important types of compounds (alkyl halides, alkenes, alkynes, aromatics, ethers, amides, mercaptans, nitriles, acid chlorides, acid anhydrides, nitro compounds, and so on), the methods introduced here can be applied equally well to other classes of compounds. The list is sufficiently broad to illustrate all the principles involved in identifying an unknown compound.

In addition, although many of the functional groups listed as being excluded will not appear as the major functional group in a compound, several of them will frequently appear as secondary, or subsidiary, functional groups. Three examples of this are presented here.



PRELIMINARY TESTS

The two combustion tests, the Beilschic test (Experiment 51B) and the ignition test (Experiment 51C) can be performed easily and quickly, and they often give valuable information. It is recommended that they be performed on all unknowns.

CHEMICAL CLASSIFICATION TESTS

The solubility tests usually suggest or eliminate several possible functional groups. The chemical classification tests listed in Experiments 51D to 51I allow you to distinguish among the possible choices. Choose only those tests that the solubility tests suggest might be meaningful. Time will be wasted performing unnecessary tests. There is no substitute for a firsthand, thorough knowledge of these tests. Study each of the sections carefully until you understand the significance of each test. Also, it will be helpful to actually try the tests on *known* substances. In this way, it will be easier to recognize a positive test. Appropriate test compounds are listed for many of the tests. When you are performing a test that is new to you, it is always good practice to run the test separately on both a known substance and the unknown *at the same time*. This practice lets you compare results directly.

Once the melting or boiling point, the solubilities, and the main chemical tests have been made, it will be possible to identify the class of compound. At this stage, with the melting point or boiling point as a guide, it will be possible to compile a list of possible compounds. Inspection of this list will suggest additional tests that must be performed to distinguish among the possibilities. For instance, one compound may be a methyl ketone and the other may not. The iodoforn test is called for to distinguish the two possibilities. The tests for the subsidiary functional groups may also be required. These tests are described in Experiments 51B and 51C. These tests should also be studied carefully; there is no substitute for firsthand knowledge about these either.

Do not perform the chemical tests either haphazardly or in a methodical, comprehensive sequence. Instead, use the tests selectively. Solubility tests automatically eliminate the need for some of the chemical tests. Each successive test will either eliminate the need for another test or dictate its use. You should also examine the tables of unknowns carefully. The boiling point or the melting point of the unknown may eliminate the need for many of the tests. For instance, the possible compounds may simply not include one with a double bond. Efficiency is the key word here. Do not waste time performing nonsensical or unnecessary tests. Many possibilities can be eliminated on the basis of logic alone.

How you proceed with the following steps may be limited by your instructor's wishes. Many instructors may restrict your access to infrared and NMR spectra until you have narrowed your choices to a few compounds *all within the same class*. Others may have you determine these data routinely. Some instructors may want students to perform elemental analysis on all unknowns; others may restrict it to only the most essential situations. Most unknowns can be identified without either spectroscopy or elemental analysis. Again, some instructors may require derivatives as a final confirmation of the compound's identity; others may not wish to use them at all.

SPECTROSCOPY

Spectroscopy is probably the most powerful and modern tool available to the chemist for determining the structure of an unknown compound. It is often possible to determine structure through spectroscopy alone. On the other hand, there are also situations for which

have corrosive vapors. Sniff any unknown substance with the greatest caution. As a first step, open the container, hold it away from you, and using your hand, carefully waft the vapors toward your nose. If you get past this stage, a closer inspection will be possible.

MELTING-POINT OR BOILING-POINT DETERMINATION

The single most useful piece of information to have for an unknown compound is its melting point or boiling point. Either piece of data will drastically limit the compounds that are possible. The electric melting-point apparatus gives a rapid and accurate measurement (see Technique 6, Section 6.7, and 6.8). To save time, you can often determine two separate melting points. The first determination can be made rapidly to get an approximate value. Then you can determine the second melting point more carefully.

The boiling point is easily obtained by a simple distillation of the unknown (Technique 8, Section 8.4), by reflux (Technique 6, Section 6.11), or by a micro boiling-point determination (Technique 6, Section 6.10). The simple distillation has the advantage in that it also purifies the compound. A Hickman head should be used if a simple distillation is performed, and you should be sure the thermometer bulb is fully immersed in the vapor of the distilling liquid. For an accurate boiling-point value, distill the liquid rapidly. You must distill more than 0.75 mL of liquid.

If the solid is high-melting ($>200^{\circ}\text{C}$), or the liquid high-boiling ($>200^{\circ}\text{C}$), a thermometer correction may be needed (Technique 6, Sections 6.12 and 6.13). In any event, allowance should be made for errors of as large as $\pm 5^{\circ}\text{C}$ in these values.

PURIFICATION

If the melting point of a solid has a wide range (ca 5°C), it should be recrystallized and the melting point redetermined.

If a liquid was highly colored before distillation, if it yielded a wide boiling-point range, or if the temperature did not hold constant during the distillation, it should be redistilled to determine a new temperature range. A reduced-pressure distillation is in order for high-boiling liquids or for those that show any sign of decomposition on heating.

Occasionally column chromatography may be necessary to purify solids that have large amounts of impurities and do not yield satisfactory results on crystallization.

Acidic or basic impurities that contaminate a neutral compound may often be removed by dissolving the compound in a low-boiling solvent, such as CH_2Cl_2 or ether, and extracting with 5% NaHCO_3 or 5% HCl , respectively. Conversely, acidic or basic compounds can be purified by dissolving them in 5% NaHCO_3 or 5% HCl , respectively, and extracting them with a low-boiling organic solvent to remove impurities. After neutralization of the aqueous solution, the desired compound can be recovered by extraction.

SOLUBILITY BEHAVIOR

Tests on solubility are described fully in Experiment 51A. They are extremely important. Determine the solubility of small amounts of the unknown in water, 5% HCl , 5% NaHCO_3 , 5% NaOH , concentrated H_2SO_4 , and organic solvents. This information reveals whether a compound is an acid, a base, or a neutral substance. The sulfuric acid test reveals whether a neutral compound has a functional group that contains an oxygen, a nitrogen, or a sulfur atom that can be protonated. This information allows you to eliminate or to choose various functional-group possibilities. The solubility tests must be made on *all* unknowns.

spectroscopy is not of much help and the traditional methods must be relied on. For this reason, you should not use spectroscopy to the exclusion of the more traditional tests but rather as a confirmation of those results. Nevertheless, the main functional groups and their immediate environmental features can be determined quickly and accurately with spectroscopy.

ELEMENTAL ANALYSIS

Elemental analysis—which allows you to determine the presence of nitrogen, sulfur, or a specific halogen atom (Cl, Br, I) in a compound—is often useful; however, other information often renders these tests unnecessary. A compound identified as an amine by solubility tests obviously contains nitrogen. Many nitrogen-containing groups (for instance, nitro groups) can be identified by infrared spectroscopy. Finally, it is not usually necessary to identify a specific halogen. The simple information that the compound contains a halogen (any halogen) may be enough information to distinguish between two compounds. A simple Beilstein test provides this information.

DERIVATIVES

One of the principal tests for the correct identification of an unknown compound comes in trying to convert the compound by a chemical reaction to another known compound. This second compound is called a **derivative**. The best derivatives are solid compounds, because the melting point of a solid provides an accurate and reliable identification of most compounds. Solids are also easily purified through crystallization. The derivative provides a way of distinguishing two otherwise very similar compounds. Usually they will have derivatives (both prepared by the same reaction) that have different melting points. Tables of unknowns and derivatives are listed in Appendix 1. Procedures for preparing derivatives are given in Appendix 2.

CONFIRMATION OF IDENTITY

A rigid and final test for identifying an unknown can be made if an “authentic” sample of the compound is available for comparison. One can compare infrared and NMR spectra of the unknown compound with the spectra of the known compound. If the spectra match, peak for peak, then the identity is probably certain. Other physical and chemical properties can also be compared. If the compound is a solid, a convenient test is the mixed melting point (Technique 6, Section 6.4). Thin-layer or gas-chromatographic comparisons may also be useful. For thin-layer analysis, however, it may be necessary to experiment with several different development solvents to reach a satisfactory conclusion about the identity of the substance in question.

Although we cannot be complete in this experiment in terms of the functional groups covered, or the tests described, the experiment should provide a good introduction to the methods and the techniques chemists use to identify unknown compounds. Textbooks that cover the subject more thoroughly are listed in the References. You are encouraged to consult these for more information, including specific methods and classification tests.

REFERENCES

- Comprehensive Textbooks**
Cheronis, N. D., and Entrikin, J. B. *Identification of Organic Compounds*. New York: Wiley-Interscience, 1963.
- Pasio, D. J., and Johnson, C. R. *Laboratory Tests for Organic Chemistry*. Englewood Cliffs, NJ: Prentice-Hall, 1979.
- Shriner, R. L., Fuson, R. C., Curtin, D. Y., and Morrill, T. C. *The Systematic Identification of Organic Compounds*, 6th ed. New York: John Wiley & Sons, 1980.
- Spectroscopy**
Bellamy, L. J. *The Infra-red Spectra of Complex Molecules*, 3rd ed. New York: Methuen, 1975.
Collup, N. B., Daly, L. H., and Wiberty, S. E. *Introduction to Infrared and Raman Spectroscopy*, 3rd ed. San Diego, CA: Academic Press, 1990.
Dyer, J. R. *Application of Absorption Spectroscopy of Organic Compounds*. Englewood Cliffs, NJ: Prentice-Hall, 1965.
Lin-Vien, D., Colthup, N. B., Failey, W. B., and Grasselli, J. G. *The Handbook of Infrared and Raman Characteristic Frequencies of Organic Molecules*. San Diego, CA: Academic Press, 1991.
Nakanishi, K. *Infrared Absorption Spectroscopy*. San Francisco: Holden-Day, 1962.
Pavia, D. L., Lampman, G. M., and Kriz, G. S., Jr. *Introduction to Spectroscopy: A Guide for Students of Organic Chemistry*, 2nd ed. Philadelphia: W. B. Saunders, 1996.
Silverstein, R. M., and Webster, F. X. *Spectrometric Identification of Organic Compounds*, 6th ed. New York: John Wiley & Sons, 1998.
- Extensive Tables of Compounds and Derivatives**
Rappoport, Z., ed. *Handbook of Tables for Organic Compound Identification*. Cleveland: Chemical Rubber Co., 1967.

Experiment 51A

Solubility Tests

Solubility tests should be performed on *every unknown*. They are extremely important in determining the nature of the main functional group of the unknown compound. The tests are very simple and require only small amounts of the unknown. In addition, solubility tests reveal whether the compound is a strong base (amine), a weak acid (phenol), a strong acid (carboxylic acid), or a neutral substance (aldehyde, ketone, alcohol, ester). The common solvents used to determine solubility types are


5% HCl	Concentrated H ₂ SO ₄
5% NaHCO ₃	Water
5% NaOH	Organic solvents

The solubility chart on page 434 indicates solvents in which compounds containing the various functional groups are likely to dissolve. The summary charts in Experiments 51D through 51I repeat this information for each functional group included in this experiment. In this section, the correct procedure for determining whether a compound is soluble in a test solvent is given. Also given is a series of explanations detailing the reasons that

compounds having specific functional groups are soluble in only specific solvents. This is accomplished by indicating the type of chemistry or the type of chemical interaction that is possible in each solvent.

WASTE DISPOSAL

Dispose of all aqueous solutions in the container designated for aqueous waste. Any remaining organic compounds must be disposed of in the appropriate organic waste container.



Solubility chart for compounds containing various functional groups.

SOLUBILITY TESTS

Procedure

Place about 1 mL of the solvent in a small test tube. Add one drop of an unknown liquid from a Pasteur pipet or a few crystals of an unknown solid from the end of a spatula, directly into the solvent. Gently tap the test tube with your finger to ensure mixing, and then observe whether any mixing lines appear in the solution. The disappearance of the liquid or solid or the appearance of

the mixing lines indicates that solution is taking place. Add several more drops of the liquid or a few more crystals of the solid to determine the extent of the compound's solubility. A common mistake in determining the solubility of a compound is testing with a quantity of the unknown too large to dissolve in the chosen solvent. Use small amounts. It may take several minutes to dissolve solids. Compounds in the form of large crystals need more time to dissolve than powders or very small crystals. In some cases, it is helpful to pulverize a compound with large crystals using a mortar and pestle. Sometimes gentle heating helps, but strong heating is discouraged, as it often leads to reaction. When colored compounds dissolve, the solution often assumes the color.

Using the preceding procedure, determine the solubility of the unknown in each of the following solvents: water, 5% HCl, 5% NaHCO₃, 5% NaOH, and concentrated H₂SO₄. With sulfuric acid, a color change may be observed rather than solution. A color change should be regarded as a positive solubility test. Solid unknowns that do not dissolve in any of the test solvents may be inorganic substances. To eliminate this possibility, determine the solubility of the unknown in several organic solvents, such as ether. If the compound is organic, a solvent that will dissolve it can usually be found.

If a compound is found to dissolve in water, the pH of the aqueous solution should be estimated with pH paper or litmus. Compounds soluble in water are usually soluble in *all* the aqueous solvents. If a compound is only slightly soluble in water, it may be *more* soluble in another aqueous solvent. For instance, a carboxylic acid may be only slightly soluble in water but very soluble in dilute base. It often will not be necessary to determine the solubility of the unknown in every solvent.

Test Compounds



Five solubility unknowns can be found on the supply shelf. The five unknowns include a base, a weak acid, a strong acid, a neutral substance with an oxygen-containing functional group, and a neutral substance that is inert. Using solubility tests, distinguish these unknowns by type. Verify your answer with the instructor.

Solubility in Water. Compounds that contain four or fewer carbons and also contain oxygen, nitrogen, or sulfur are often soluble in water. Almost any functional group containing these elements will lead to water solubility for low-molecular-weight (C₄) compounds. Compounds having five or six carbons and any of those elements are often insoluble in water or have borderline solubility. Branching of the alkyl chain in a compound lowers the intermolecular forces between its molecules. This is usually reflected in a lowered boiling point or melting point and a greater solubility in water for the branched compound than for the corresponding straight-chain compound. This occurs simply because the molecules of the branched compound are more easily separated from one another. Thus, *t*-butyl alcohol would be expected to be more soluble in water than *n*-butyl alcohol.


When the ratio of the oxygen, nitrogen, or sulfur atoms in a compound to the carbon atoms is increased, the solubility of that compound in water often increases. This is due to the increased number of polar functional groups. Thus, 1,5-pentanediol would be expected to be more soluble in water than 1-pentanol.

As the size of the alkyl chain of a compound is increased beyond about four carbons, the influence of a polar functional group is diminished, and the water solubility begins to decrease. A few examples of these generalizations are given here.

Compounds dissolve in base because they form sodium salts that are soluble in the aqueous medium. The salts of some high-molecular-weight compounds are not soluble, however, and precipitate. The salts of the long-chain carboxylic acids, such as myristic acid C_{14} , palmitic acid C_{16} , and stearic acid C_{18} , which form soaps, are in this category. Some phenols also produce insoluble sodium salts, and often these are colored due to resonance in the anion.


Strong Acids (soluble in both NaOH and NaHCO_3)	Weak Acids (soluble in NaOH but not NaHCO_3)
Sulfonic acids RSO_3H Carboxylic acids RCOOH <i>Ortho</i> - and <i>para</i> -substituted di- and trinitrophenols 	Phenols Nitroalkanes β -Diketones β -Diesters Imides Sulfonamides 

Both phenols and carboxylic acids produce resonance-stabilized conjugate bases. Thus, bases of the appropriate strength may easily remove their acidic protons to form the sodium salts.



In phenols, substitution of nitro groups in the *ortho* and *para* positions of the ring increases the acidity. Nitro groups in these positions provide additional delocalization in the conjugate anion. Phenols that have two or three nitro groups in the *ortho* and *para* positions often dissolve in both sodium hydroxide and sodium bicarbonate solutions.


Solubility in Concentrated Sulfuric Acid. Many compounds are soluble in cold concentrated sulfuric acid. Of the compounds included in this experiment, alcohols, ketones, aldehydes, and esters are in this category. Other compounds that also dissolve include



Solubility in 5% HCl. The possibility of an amine should be considered immediately if a compound is soluble in dilute acid (5% HCl). Aliphatic amines (RNH_2 , R_2NH , R_3N) are basic compounds that readily dissolve in acid because they form hydrochloride salts that are soluble in the aqueous medium:



The substitution of an aromatic (benzene) ring Ar for an alkyl group R reduces the basicity of an amine somewhat, but the amine will still protonate, and it will still generally be soluble in dilute acid. The reduction in basicity in an aromatic amine is due to the resonance delocalization of the unshared electrons on the amino nitrogen of the free base. The delocalization is lost on protonation, a problem that does not exist for aliphatic amines. The substitution of two or three aromatic rings on an amine nitrogen reduces the basicity of the amine even further. Diaryl and triaryl amines do not dissolve in dilute HCl because they do not protonate easily. Thus, Ar_2NH and Ar_3N are insoluble in dilute acid. Some amines of very high molecular weight, such as tribromoamine ($\text{MW} = 330$), may also be insoluble in dilute acid.






Solubility in 5% NaHCO_3 and 5% NaOH. Compounds that dissolve in sodium bicarbonate, a weak base, are strong acids. Compounds that dissolve in sodium hydroxide, a strong base, may be either strong or weak acids. Thus, one can distinguish weak and strong acids by determining their solubility in both strong (NaOH) and weak (NaHCO_3) base. The classification of some functional groups as either weak or strong acids is given in the table on page 437.

In this experiment, carboxylic acids ($\text{pK}_a \sim 5$) are generally indicated when a compound is soluble in both bases, while phenols ($\text{pK}_a \sim 10$) are indicated when it is soluble in NaOH only.

alkenes, alkynes, ethers, nitroaromatics, and amides. Because several different kinds of compounds are soluble in sulfuric acid, further chemical tests and spectroscopy will be needed to differentiate among them.


Compounds that are soluble in concentrated sulfuric acid but not in dilute acid are extremely weak bases. Almost any compound containing a nitrogen, an oxygen, or a sulfur atom can be protonated in concentrated sulfuric acid. The ions produced are soluble in the medium.



Inert Compounds. Compounds not soluble in concentrated sulfuric acid or any of the other solvents are said to be **inert**. Compounds not soluble in concentrated sulfuric acid include the alkanes, most simple aromatics, and the alkyl halides. Some examples of inert compounds are hexane, benzene, chlorobenzene, chlorohexane, and toluene.

Experiment 51B

Tests for the Elements (N, S, X)



Except for amines (Experiment 51C), which are easily detected by their solubility behavior, all compounds issued in this experiment will contain heteroelements (N, S, Cl, Br, or I) only as *secondary* functional group. These will be subsidiary to some other important functional group. Thus, no alkyl or aryl halides, nitro compounds, thiols, or thioethers will be issued. However, some of the unknowns may contain a halogen or a nitro group. Less frequently, they may contain a sulfur atom or a cyano group.

Consider as an example *p*-bromobenzaldehyde, an aldehyde that contains bromine as a ring substituent. The identification of this compound would hinge on whether the investigator could identify it as an aldehyde. It could probably be identified *without* proving the existence of bromine in the molecule. That information, however, could make the identification easier. In this experiment, methods are given for identifying the presence of a halogen or a nitro group in an unknown compound. Also given is a general method (sodium fusion) for detecting the principal heteroelements that may exist in organic molecules.

Classification Tests

Halides	Nitro Groups	N, S, X (Cl, Br, I)
Beilstein test Silver nitrate Sodium iodide/acetone	Ferrous hydroxide	Sodium fusion

WASTE DISPOSAL

Dispose of all solutions containing silver into a waste container designated for this purpose. Any other aqueous solutions should be disposed of in the container designated for aqueous waste. Any remaining organic compounds must be disposed of in the appropriate organic waste container under the hood. This is particularly true of any solution containing benzyl bromide, which is a lachrymator.

TESTS FOR A HALIDE

Beilstein Test

Procedure. Bend a small loop in the end of a short length of copper wire. Heat the loop end of the wire in a Bunsen burner flame. After cooling, dip the wire directly into a small sample of the unknown. If the unknown is a solid and won't adhere to the copper wire, place a small amount of the substance on a watch glass, wet the copper wire in distilled water, and place the wire into the sample on the watch glass. The solid should adhere to the wire. Now heat the wire in the Bunsen burner flame again. The compound will first burn. After the burning, a green flame will be produced if a halogen is present.

Test Compounds. Try this test on bromobenzene and benzoic acid.

Halogens can be detected easily and reliably by the Beilstein test. It is the simplest method for determining the presence of a halogen, but it does not differentiate among chlorine, bromine, and iodine, any one of which will give a positive test. However, when the identity of the unknown has been narrowed to two choices, of which one has a halogen and one does not, the Beilstein test will often be enough to distinguish between the two.

A positive Beilstein test results from the production of a volatile copper halide when an organic halide is heated with copper oxide. The copper halide imparts a blue-green color to the flame.


This test can be very sensitive to small amounts of halide impurities in some compounds. Therefore, use caution in interpreting the results of the test if you obtain only a weak color.

Silver Nitrate Test

Procedure. Add one drop of a liquid or five drops of a concentrated ethanolic solution of a solid unknown to 2 mL of a 2% ethanolic silver nitrate solution. If no reaction is observed after 5 minutes at room temperature, heat the solution in a hot water bath at about 100°C and note whether a precipitate forms. If a precipitate forms, add two drops of 5% nitric acid and note whether the precipitate dissolves. Carboxylic acids give a false test by precipitating in silver nitrate, but they dissolve when nitric acid is added. Silver halides, in contrast, do not dissolve in nitric acid.


Test Compounds. Apply this test to benzyl bromide (α -bromotoluene) and bromobenzene. Discard all waste reagents in a suitable waste container in the hood, because benzyl bromide is a lachrymator.

This test depends on the formation of a white or off-white precipitate of silver halide when silver nitrate is allowed to react with a sufficiently reactive halide.



The test does not distinguish among chlorides, bromides, and iodides but does distinguish labile (reactive) halides from halides that are unreactive. Halides substituted on an aromatic ring will not usually give a positive silver nitrate test; however, alkyl halides of many types will give a positive test.

The most reactive compounds are those able to form stable carbocations in solution and those equipped with good leaving groups (X = I, Br, Cl). Benzyl, allyl, and tertiary halides react immediately with silver nitrate. Secondary and primary halides do not react at room temperature but react readily when heated. Aryl and vinyl halides do not react at all, even at elevated temperatures. This pattern of reactivity fits the stability order for various carbocations quite well. Compounds that produce stable carbocations react at higher rates than those that do not.



The fast reaction of benzylic and allylic halides is a result of the resonance stabilization that is available to the intermediate carbocations formed. Tertiary halides are more reactive than secondary halides, which are in turn more reactive than primary or methyl halides because alkyl substituents are able to stabilize the intermediate carbocations by an electron-releasing effect. The methyl carbocations have no alkyl groups and are the least stable of all carbocations mentioned thus far. Vinyl and aryl carbocations are extremely unstable because the charge is localized on an sp^2 -hybridized carbon (double-bond carbon) rather than on one that is sp^3 -hybridized.

Sodium Iodide in Acetone

Procedure. This test is described in Experiment 18.

DETECTION OF NITRO GROUPS


Although nitro compounds will not be issued as distinct unknowns, many of the unknowns may have a nitro group as a secondary functional group. The presence of a nitro group, and hence nitrogen, in an unknown compound is determined most easily by infrared spectroscopy. However, many nitro compounds give a positive result in the following test. Unfortunately, functional groups other than the nitro group may also give a positive result. You should interpret the results of this test with caution.

Ferrous Hydroxide Test

Procedure. Place 1.5 mL of freshly prepared 5% aqueous ferrous ammonium sulfate in a small test tube, and add about 10 mg of the unknown compound. Mix the solution well, and then add first one drop of 2M sulfuric acid, and then 1 mL of 2M potassium hydroxide in methanol. Stopper the test tube, and shake it vigorously. A positive test is indicated by the formation of a red-brown precipitate, usually within 1 minute.

Test Compound. 2-Nitrotoluene

Most nitro compounds oxidize ferrous hydroxide to ferric hydroxide, which is a red-brown solid. A precipitate indicates a positive test.




Infrared Spectroscopy

The nitro group gives two strong bands near 1560 and 1350 cm^{-1} .

DETECTION OF A CYANO GROUP


Although nitriles will not be given as unknowns in this experiment, the cyano group may be a subsidiary functional group whose presence or absence is important to the final identification of an unknown compound. The cyano group can be hydrolyzed in a strong base, by heating vigorously, to give a carboxylic acid and ammonia gas:



The ammonia can be detected by its odor or by moist pH paper. However, this method is somewhat difficult, and the presence of a nitrile group is confirmed most easily by infrared spectroscopy. No other functional groups (except some $\text{C}\equiv\text{C}$) absorb in the same region of the spectrum as $\text{C}\equiv\text{N}$.

Experiment 51C

Tests for Unsaturation



The unknowns to be issued for this experiment have neither a double bond nor a triple bond as their *only* functional group. Hence, simple alkenes and alkynes can be ruled out as possible compounds. Some of the unknowns may have a double or a triple bond, however *in addition* to another more important functional group. The tests described allow you to determine the presence of a double bond or a triple bond (unsaturation) in such compounds.

Classification Tests

Unsaturation	Aromaticity
Bromine-carbon tetrachloride Potassium permanganate	Ignition test

WASTE DISPOSAL

Test reagents that contain bromine should be discarded into a special waste container designated for this purpose. Methylene chloride and carbon tetrachloride must be placed in the organic waste container designated for the disposal of halogenated organic wastes. Dispose of all other aqueous solutions in the container designated for aqueous waste. Any remaining organic compounds must be disposed of in the appropriate organic waste container.

TESTS FOR SIMPLE MULTIPLE BONDS


Bromine in Carbon Tetrachloride or Methylene Chloride

Procedure. Dissolve 50 mg of a solid unknown or two drops of a liquid unknown in 1 mL of carbon tetrachloride (or 1,2-dimethoxyethane). Add a 2% (by volume) solution of bromine in carbon tetrachloride, dropwise, and shake it until the bromine color persists. The test is positive if more than five drops of the bromine solution are needed so that the color remains for 1 minute. Usually, many drops of the bromine solution will be needed if unsaturation is present. Hydrogen bromide should not be evolved. If hydrogen bromide gas is evolved, you will note a "fog" while you blow across the mouth of the test tube. The HBr can also be detected by a moistened piece of litmus or pH paper. If hydrogen bromide is evolved, the reaction is a **substitution reaction** (see discussion below) and not an **addition reaction**, and a double or triple bond is probably not present.


Methylene Chloride. Even though carbon tetrachloride is used in very small quantities in this test, it poses certain health hazards (see p. 17), and another solvent may be preferable. Methylene chloride (dichloromethane) can be substituted for carbon tetrachloride. Certain problems arise, however, because methylene chloride slowly reacts with bromine, presumably by a light-induced free-radical process, to produce HBr. After about 1 week, the color of a 2% solution of bromine in methylene chloride fades noticeably, and the odor of the HBr can be detected in the reagent. Although the decolorization tests still work satisfactorily, the presence of HBr makes it difficult to distinguish between addition and substitution reactions. A freshly prepared solution of bromine in methylene chloride must be used to make this distinction. Deterioration of the reagent can be forestalled by storing it in a brown glass bottle. Most other substitute solvents also present problems. Ethers, for instance, react slowly in the same way as methylene chloride, and hydrocarbons, such as hexane, are not general enough solvents to be able to dissolve all the possible test compounds.

Test Compounds. Try this test with cyclohexene, cyclohexane, toluene, and acetone.


A successful test depends on the addition of bromine, a red liquid, to a double or a triple bond to give a colorless dibromide:



Not all double bonds react with bromine-carbon tetrachloride solution. Only those that are electron-rich are sufficiently reactive nucleophiles to initiate the reaction. A double bond that is substituted by electron-withdrawing groups often fails to react or reacts slowly. Fumaric acid is an example of a compound that fails to give the reaction.



Aromatic compounds either do not react with bromine-carbon tetrachloride reagent or they react by **substitution**. Only the aromatic rings that have activating groups as substituents ($-\text{OH}$, $-\text{OR}$, or $-\text{NR}_2$) give the substitution reaction.




Some ketones and aldehydes react with bromine to give a **substitution product**, but this reaction is slow except for ketones that have a high enol content. When substitution occurs, not only is the bromine color discharged, but hydrogen bromide gas is also evolved.

Potassium Permanganate (Baeyer Test)

Procedure. Dissolve 25 mg of a solid unknown or two drops of a liquid unknown in 2 mL of water or 95% ethanol (1,2-dimethoxyethane may also be used). Slowly add a 1% aqueous solution (weight/volume) of potassium permanganate, drop by drop while shaking, to the unknown. In a positive test, the purple color of the reagent is discharged, and a brown precipitate of manganese dioxide forms, usually within 1 minute. If alcohol was the solvent, the solution should not be allowed to stand for more than 5 minutes, because oxidation of the alcohol will begin slowly. Because permanganate solutions undergo some decomposition to manganese dioxide on standing, any small amount of amount of precipitate should be interpreted with caution.

Test Compounds. Try this test on cyclohexene and toluene.

This test is positive for double and triple bonds but not for aromatic rings. It depends on the conversion of the purple ion MnO_4^- to a brown precipitate of MnO_2 following the oxidation of an unsaturated compound.



Other easily oxidized compounds also give a positive test with potassium permanganate solution. These substances include aldehydes, some alcohols, phenols, and aromatic amines. If you suspect that any of these functional groups are present, you should interpret the test with caution.

Spectroscopy

Infrared

Double Bonds (C=C)

C=C stretch usually occurs near 1680–1620 cm^{-1} . Symmetrical alkenes may have no absorption.

C—H stretch of vinyl hydrogens occurs $>3000 \text{ cm}^{-1}$, but usually not higher than 3150 cm^{-1} .

C—H out-of-plane bending occurs near 1000–700 cm^{-1} (see Appendix 3).

Triple Bonds (C≡C)

C≡C stretch usually occurs near 2250–2100 cm^{-1} . The peak is usually sharp. Symmetrical alkynes show

no absorption.

C—H stretch of terminal acetylenes occurs near 3310–3200 cm^{-1} .

TESTS FOR AROMATICITY

None of the unknowns to be issued for this experiment will be simple aromatic hydrocarbons. All aromatic compounds will have a principal functional group as a part of their structure. Nevertheless, in many cases it will be useful to be able to recognize the presence of an aromatic ring. Although spectroscopy provides the easiest method of determining aromatic systems, often they can be detected by a simple ignition test.

Ignition Test

Procedure. Working in a hood, place a small amount of the compound on a spatula, and place it in the flame of a Bunsen burner. Observe whether a sooty flame is the result. Compounds giving a sooty yellow flame have a high degree of unsaturation and may be aromatic.

Test Compound. Try this test with naphthalene.

The presence of an aromatic ring or other centers of unsaturation will lead to the production of a sooty yellow flame in this test. Compounds that contain little oxygen and have a high carbon-to-hydrogen ratio, burn at a low temperature with a yellow flame. Much carbon is produced when they are burned. Compounds that contain oxygen generally burn at a higher temperature with a clean blue flame.

Spectroscopy

Infrared

C=C aromatic ring double bonds appear in the 1600–1450 cm^{-1} region. There are often four sharp absorptions that occur in pairs near 1600 cm^{-1} and 1450 cm^{-1} which are characteristic of an aromatic ring.

Special ring absorptions: There are often weak ring absorptions around 2000–1600 cm^{-1} . These are often obscured, but when they can be observed, the relative shapes and numbers of these peaks can often be used to ascertain the type of ring substitution (see Appendix 3).


=C—H stretch, aromatic ring: The aromatic C—H stretch always occurs at a higher frequency than 3000 cm^{-1} .

=C—H out-of-plane bending peaks appear in the region 900–690 cm^{-1} . The number and position of these peaks can be used to determine the substitution pattern of the ring (see Appendix 3).

Nuclear Magnetic Resonance

Hydrogens attached to an aromatic ring usually have resonance near 7 ppm. Monosubstituted rings not substituted by anisotropic or electronegative groups usually give a single resonance for all the ring hydrogens. Monosubstituted rings with anisotropic or electronegative groups usually have the aromatic resonances split into two groups integrating either 3:2 or 2:3. A nonsymmetric, *para*-disubstituted ring has a characteristic four-peak splitting pattern (see Appendix 4).

Experiment 51D Aldehydes and Ketones



Compounds containing the carbonyl functional group $\text{C}=\text{O}$, where it has only hydrogen atoms or alkyl groups as substituents, are called aldehydes RCHO or ketones RCOR . The chemistry of these compounds is primarily due to the chemistry of the carbonyl functional groups. These compounds are identified by the distinctive reactions of the carbonyl function.

Solubility Characteristics	Classification Tests
HCl NaHCO_3 NaOH H_2SO_4 Ether (-) (-) (-) (+) (+)	Aldehydes and ketones 2,4-Dinitrophenylhydrazine
Water: $<C_7$ and some $C_8(+)$ $>C_8(-)$	Aldehydes only Methyl ketones Chromic acid Iodoform test Tollens reagent Compounds with high enol content Ferric chloride test

WASTE DISPOSAL

Solutions containing 2,4-dinitrophenylhydrazine or derivatives formed from it should be placed in a waste container designated for these compounds. Any solution containing chromium must be disposed of in a waste container specifically identified for the disposal of chromium wastes. Dispose of all solutions containing silver by acidifying them with 5% hydrochloric acid and then placing them in a waste container designated for this purpose. Dispose of all other aqueous solutions in the container designated for aqueous waste. Any remaining organic compounds must be disposed of in the appropriate organic waste container.

CLASSIFICATION TESTS

Most aldehydes and ketones give a solid, yellow-to-red precipitate when mixed with 2,4-dinitrophenylhydrazine. However, only aldehydes will reduce chromium(VI) or silver(I). By this difference in behavior, you can differentiate between aldehydes and ketones.

2,4-Dinitrophenylhydrazine


Procedure. Place one drop of the liquid unknown in a small test tube, and add 1 mL of the 2,4-dinitrophenylhydrazine reagent. If the unknown is a solid, dissolve about 10 mg (estimate) in

a minimum amount of 95% ethanol or bis(2-ethoxyethyl) ether before adding the reagent. Shake the mixture vigorously. Most aldehydes and ketones will give a yellow-to-red precipitate immediately. However, some compounds will require up to 15 minutes, or even *gentle* heating, to give a precipitate. A precipitate indicates a positive test.

Test Compounds. Try this test on cyclohexanone, benzaldehyde, and benzophenone.

CAUTION: Many derivatives of phenylhydrazine are suspected carcinogens (see p. 19) and should be handled with care. Avoid contact.

Reagent. Dissolve 3.0 g of 2,4-dinitrophenylhydrazine in 15 mL of concentrated sulfuric acid. In a beaker mix 20 mL of water and 70 mL of 95% ethanol. While stirring vigorously, add the 2,4-dinitrophenylhydrazine solution to the aqueous ethanol mixture slowly. After thorough mixing, filter the solution by gravity through a fluted filter.



Most aldehydes and ketones give a precipitate, but esters generally do not give this result. Thus, an ester usually can be eliminated by this test. The color of the 2,4-dinitrophenylhydrazone (precipitate) formed is often a guide to the amount of conjugation in the original aldehyde or ketone. Unconjugated ketones, such as cyclohexanone, give yellow precipitates, whereas conjugated ketones, such as benzophenone, give orange-to-red precipitates. Compounds that are highly conjugated give red precipitates. However, the 2,4-dinitrophenylhydrazine reagent is itself orange-red, and the color of any precipitate must be judged cautiously. Occasionally compounds that are either strongly basic or strongly acidic precipitate the unreacted reagent.

Some allylic and benzylic alcohols give this test result because the reagent can oxidize them to aldehydes and ketones, which subsequently react. Some alcohols may be contaminated with carbonyl impurities, either because of their method of synthesis (reduction) or because they have become air-oxidized. A precipitate formed from small amounts of impurity in the solution will be formed in small amount. With some caution, a test that gives only a slight amount of precipitate can usually be ignored. The infrared spectrum of the compound should establish its identity and identify any impurities present.

Chromic Acid Test

Procedure. Dissolve one drop of a liquid or 10 mg (approximate) of a solid aldehyde in 1 mL of reagent-grade acetone. Add several drops of the chromic acid reagent, a drop at a time while shaking the mixture. A positive test is indicated by a green precipitate and a loss of the orange

color in the reagent. With aliphatic aldehydes RCHO, the solution turns cloudy within 5 seconds and a precipitate appears within 30 seconds. With aromatic aldehydes ArCHO, it generally takes 30–120 seconds for a precipitate to form, but with some it may take even longer.

In a negative test, there is usually no precipitate. In some cases, however, a precipitate forms, but the solution remains orange.


In performing this test, make quite sure that the acetone used for the solvent does not give a positive test with the reagent. Add several drops of the chromic acid reagent to a few drops of the reagent acetone contained in a small test tube. Allow this mixture to stand for 3–5 minutes. If no reaction has occurred by this time, the acetone is pure enough to use as a solvent for the test. If a positive test resulted, try another bottle of acetone, or distill some acetone from potassium permanganate to purify it.

Test Compounds. Try this test on benzaldehyde, butanal (butyraldehyde), and cyclonexanone.

CAUTION: Many compounds of chromium (VI) are suspected carcinogens (see p. 19) and should be handled with care. Avoid contact.

Reagent. Dissolve 1.0 g of chromic oxide CrO₃ in 1 mL of concentrated sulfuric acid. Then dilute this mixture carefully with 3 mL of water.

This test has as its basis the fact that aldehydes are easily oxidized to the corresponding carboxylic acid by chromic acid. The green precipitate is due to chromous sulfate.



Primary and secondary alcohols are also oxidized by this reagent (see Experiment 51H). Therefore, this test is not useful in identifying aldehydes *unless* a positive identification of the carbonyl group has already been made. Aldehydes give a 2,4-dinitrophenylhydrazine test result, whereas alcohols do not.

There are numerous other tests used to detect the aldehyde functional group. Most are based on an easily detectable oxidation of the aldehyde to a carboxylic acid. The most common tests are the Tollens, Fehling, and Benedict tests. Only the Tollens test is described here.

Tollens Test

Procedure. The reagent must be prepared immediately before use. To prepare the reagent, mix 1 mL of Tollens solution A with 1 mL of Tollens solution B. A precipitate of silver oxide will form. Add enough dilute (10%) ammonia solution (dropwise) to the mixture to dissolve the silver oxide *just barely*. The reagent so prepared can be used immediately for the following test.


Dissolve one drop of a liquid aldehyde or 10 mg (approximate) of a solid aldehyde in the minimum amount of bis(2-ethoxyethyl) ether. Add this solution, a little at a time, to the 2–3 mL of reagent contained in a small test tube. Shake the solution well. If a mirror of silver is deposited on the inner walls of the test tube, the test is positive. In some cases, it may be necessary to warm the test tube in a bath of warm water.

CAUTION: The reagent should be prepared immediately before use and all residues disposed of immediately after use. Dispose of any residues by acidifying them with 5% hydrochloric acid and then placing them in a waste container designated for this purpose. On standing, the reagent tends to form silver fulminate, a very explosive substance. Solutions containing the mixed Tollens reagent should never be stored.

Test Compounds. Try the test on acetone and benzaldehyde.

Reagents. *Solution A:* Dissolve 3.0 g of silver nitrate in 30 mL of water. *Solution B:* Prepare a 10% sodium hydroxide solution.

Most aldehydes reduce ammoniacal silver nitrate solution to give a precipitate of silver metal. The aldehyde is oxidized to a carboxylic acid:



Ordinary ketones do not give a positive result in this test. The test should be used only if it has already been shown that the unknown compound is either an aldehyde or a ketone.

Iodoform Test

Procedure. Prepare a 60 to 70°C water bath in a beaker. Using a Pasteur pipet, add six drops of a liquid unknown to a 15 × 100-mm or 15 × 125-mm test tube. Alternatively, 0.06 g of a solid unknown may be used. Dissolve the liquid or solid unknown compound in 2 mL of 1,2-dimethoxyethane. Add 2 mL of 10% aqueous sodium hydroxide solution, and place the test tube in the hot water bath. Next add 4 mL of iodine–potassium iodide solution in 1-mL portions to the test tube. Cork the test tube, and shake it after adding each portion of iodine reagent. Heat the mixture in the hot water bath for about 5 minutes, shaking the test tube occasionally. It is likely that some or all the dark color of the iodine reagent will be discharged.

If the dark color of the iodine reagent is still apparent following heating, add 10% sodium hydroxide solution until the dark color of the iodine reagent has been discharged. Shake the mixture in the test tube (corked) during the addition of the sodium hydroxide. Care need not be taken to avoid adding excess sodium hydroxide.


After the dark iodine color of the solution has been discharged, fill the test tube with water to within 2 cm of the top. Cork the test tube, and shake it vigorously. Allow the tube to stand for at least 15 minutes at room temperature. The appearance of a pale yellow precipitate of iodoform CHI₃ constitutes a positive test, indicating that the unknown is a methyl ketone or a compound that is easily oxidized to a methyl ketone, such as a 2-alkanone. Other ketones will also decolorize the iodine solution, but they will not give a precipitate of iodoform *unless* there is an impurity of a methyl ketone present in the unknown.

The yellow precipitate usually settles out slowly onto the bottom of the test tube. Sometimes the yellow color of iodoform is masked by a dark substance. If this is the case, cork the test tube and shake it vigorously. If the dark color persists, add more sodium hydroxide solution and shake the test tube again. Then allow the tube to stand for at least 15 minutes. If there is some doubt as to whether the solid is iodoform, collect the precipitate on a Hirsch funnel and dry it. Iodoform melts at 119–121°C.

Test Compounds. Try the test on 2-heptanone, 4-heptanone (dipropyl ketone), and 2-pentanone.

Reagents. The iodine reagent is prepared by dissolving 20 g of potassium iodide and 10 g of iodine in 100 mL of water. The aqueous sodium hydroxide solution is prepared by dissolving 10 g of sodium hydroxide in 100 mL of water.

The basis of this test is the ability of certain compounds to form a precipitate of iodoform when treated with a basic solution of iodine. Methyl ketones are the most common types of compounds that give a positive result in this test. However, acetaldehyde CH_3CHO and alcohols with the hydroxyl group at the 2-position of the chain also give a precipitate of iodoform. 2-Alkanols of the type described are easily oxidized to methyl ketones under the conditions of the reaction. The other product of the reaction, besides iodoform, is the sodium or potassium salt of a carboxylic acid.



Ferric Chloride Test

Procedure. Some aldehydes and ketones, those that have a high enol content, give a positive ferric chloride test, as described for phenols in Experiment 51F.

Spectroscopy

Infrared

The carbonyl group is usually one of the strongest-absorbing groups in the infrared spectrum, with a very broad range: $1800\text{--}1650\text{ cm}^{-1}$. The aldehyde functional group has very characteristic C—H stretch absorptions: two sharp peaks that lie *far outside* the usual region for $-\text{C}-\text{H}$, $=\text{C}-\text{H}$, or $\equiv\text{C}-\text{H}$.

Aldehydes

C=O stretch at approximately 1725 cm^{-1} is normal. $1725\text{--}1685\text{ cm}^{-1,2}$

C—H stretch (aldehyde—CHO) has two weak bands at about 2750 cm^{-1} and 2850 cm^{-1} .

Ketones

C=O stretch at approximately 1715 cm^{-1} is normal. $1780\text{--}1665\text{ cm}^{-1,2}$


Nuclear Magnetic Resonance

Hydrogens alpha to a carbonyl group have resonance in the region between 2 and 3 ppm. The hydrogen of an aldehyde group has a characteristic resonance between 9 and 10 ppm. In aldehydes, there is coupling between the aldehyde hydrogen and any alpha hydrogens ($J = 1\text{--}3\text{ Hz}$).

²Conjugation moves the absorption to lower frequencies. Ring strains (cyclic ketones) moves the absorption to higher frequencies.

Derivatives

The most common derivatives of aldehydes and ketones are 2,4-dinitrophenylhydrazones, oximes, and semicarbazones. Procedures for preparing these derivatives are given in Appendix 2.



Experiment 51E Carboxylic Acids



Carboxylic acids are detectable mainly by their solubility characteristics. They are soluble in both dilute sodium hydroxide and sodium bicarbonate solutions.

Solubility Characteristics	Classification Tests
HCl (-)	pH of an aqueous solution
NaHCO ₃ (+)	Sodium bicarbonate
NaOH (+)	Silver nitrate
H ₂ SO ₄ (+)	Neutralization equivalent
Water: <C ₆ (+) >C ₆ (-)	

WASTE DISPOSAL

Dispose of all aqueous solutions in the container designed for aqueous waste. Any remaining organic compounds must be disposed of in the appropriate organic waste container.

Appendix 1

Tables of Unknowns and Derivatives

More extensive tables of unknowns may be found in Z. Rappoport, ed. *Handbook of Tables for Organic Compound Identification*, 3rd ed. Cleveland: Chemical Rubber Co., 1967.

Aldehydes

Compound	bp	mp	Semi-carbazone*	2,4-Dinitro-phenyl-hydrazone*
Ethanal (acetaldehyde)	21	—	162	168
Propanal (propionaldehyde)	48	—	89	148
Propional (acrolein)	52	—	171	165
2-Methylpropanal (isobutyraldehyde)	64	—	125	187
Butanal (butyraldehyde)	75	—	95	123
3-Methylbutanal (isovaleraldehyde)	92	—	107	123
Pentanal (valeraldehyde)	102	—	—	106
2-Butenal (crotonaldehyde)	104	—	199	190
2-Ethylbutanal (diethylacetaldehyde)	117	—	99	95
Hexanal (caproaldehyde)	130	—	106	104
Heptanal (heptaldehyde)	153	—	109	108
2-Furaldehyde (furfural)	162	—	202	212
2-Ethylhexanal	163	—	254	114
Octanal (caprylaldehyde)	171	—	101	106
Benzaldehyde	179	—	222	237
Phenylethanal (phenylacetaldehyde)	195	33	153	121
2-Hydroxybenzaldehyde (salicylaldehyde)	197	—	231	248
4-Methylbenzaldehyde (<i>p</i> -tolualdehyde)	204	—	234	234
3,7-Dimethyl-6-octenal (citronellal)	207	—	82	77
2-Chlorobenzaldehyde	213	11	229	213
4-Methoxybenzaldehyde (<i>p</i> -anisaldehyde)	248	2.5	210	253
<i>trans</i> -Cinnamaldehyde	250 d.	—	215	255
3,4-Methylenedioxybenzaldehyde (piperonal)	263	37	230	266 d.
2-Methoxybenzaldehyde (<i>o</i> -anisaldehyde)	245	38	215 d.	254
4-Chlorobenzaldehyde	214	48	230	254
3-Nitrobenzaldehyde	—	58	246	293
4-Dimethylaminobenzaldehyde	—	74	222	325
Vanillin	285 d.	82	230	271
4-Nitrobenzaldehyde	—	106	221	320 d.
4-Hydroxybenzaldehyde	—	116	224	280 d.
(±)-Glyceraldehyde	—	142	160 d.	167

Note: "d." indicates "decomposition."

*See Appendix 2, "Procedures for Preparing Derivatives."

Ketones

Compound	bp	mp	Semi-carbazone*	2,4-Dinitro-phenyl-hydrazone*
2-Propanone (acetone)	56	—	187	126
2-Butanone (methyl ethyl ketone)	80	—	146	117
3-Methyl-2-butanone (isopropyl methyl ketone)	94	—	112	120
2-Pentanone (methyl propyl ketone)	101	—	112	143
3-Pentanone (diethyl ketone)	102	—	138	156
Pinacolone	106	—	157	125
4-Methyl-2-pentanone (isobutyl methyl ketone)	117	—	132	95
2,4-Dimethyl-3-pentanone (diisopropyl ketone)	124	—	160	95
2-Hexanone (methyl butyl ketone)	128	—	125	106
4-Methyl-3-penten-2-one (mesityl oxide)	130	—	164	205
Cyclohexanone	131	—	210	146
2,3-Pentanedione	134	—	122 (mono) 209 (di)	209
2,4-Pentanedione (acetylacetone)	139	—	—	122 (mono) 209 (di)
4-Heptanone (dipropyl ketone)	144	—	132	75
2-Heptanone (methyl amyl ketone)	151	—	123	89
Cyclohexanone	156	—	166	162
2,6-Dimethyl-4-heptanone (diisobutyl ketone)	168	—	122	92
2-Octanone	173	—	122	58
Cycloheptanone	181	—	163	148
2,5-Hexanedione (acetylacetone)	191	-9	185 (mono) 224 (di)	257 (di)
Acetophenone (methyl phenyl ketone)	202	20	198	238
Phenyl-2-propanone (phenylacetone)	216	27	198	156
Propiophenone (ethyl phenyl ketone)	218	21	182	191
4-Methylacetophenone	226	—	205	258
2-Undecanone	231	12	122	63
4-Chloroacetophenone	232	12	204	236
4-Phenyl-2-butanone (benzylacetone)	235	—	142	127
4-Chloropropiophenone	—	36	176	223
4-Phenyl-3-buten-2-one	37	37	187	227
4-Methoxyacetophenone	258	38	198	228
Benzophenone	305	48	167	238
4-Bromoacetophenone	225	51	208	230
2-Acetonaphthone	—	54	235	262
Desoxybenzoin	320	60	148	204
3-Nitroacetophenone	202	80	257	228
9-Fluorenone	345	83	234	283
Benzoin	344	136	206	245
4-Hydroxypropiophenone	—	148	—	229
(±)-Camphor	205	179	237	177

*See Appendix 2, "Procedures for Preparing Derivatives."